



NYS Mental Health Education in Schools Law

Summary and Recommendations

Who	In NYS, health education is a K-12 requirement; therefore, all students in public and private schools, grades K-12 are learning about mental health.
What	<p>The law takes a holistic view of mental health ensuring that students “recognize the multiple dimensions of health by including mental health, and the relationship of physical health and mental health, so as to enhance student understanding, attitudes and behaviors that promote health, well-being and human dignity.”</p> <ul style="list-style-type: none"> • The NYS Education Department released a framework for Mental Health Education Instruction aligned with the NYS Health Education Standards as part of <i>Mental Health Education Literacy in Schools: Linking to a Continuum of Well-Being</i> (pages 13-24).
Where	While the law specifically pertains to the health education program, many schools have taken the opportunity to integrate mental health as a topic for discussion in other subjects, such as ELA, Participation in Government, History and Biology.
When	Considering that students only receive two semesters of Health between 7 th and 12 th grades, schools should consider developing an on-going, integrated approach to raising awareness about mental health. Some schools have established student-led wellness clubs that organize resource fairs, draft social media posts, coordinate guest speakers and facilitate mental health and wellness activities for younger students. Other schools have developed school-wide awareness campaigns, such as “Wellness Wednesdays”, using morning announcements to share health and safety information, such as stress management strategies, appropriate web-based resources, and the importance of sleep, staying hydrated or the use of sunscreen.
Why	Research suggests that an increase in mental health literacy leads to early intervention and better health outcomes. According to the National Institute of Mental Health, nearly half of all youth will experience a mental health challenge before the age of 18. We all have mental health and we all benefit from greater understanding of how we can support our own mental health and wellness and that of others.
How	The World Health Organization recommends a public health approach to mental health; a multi-tiered approach that includes the whole school community at the universal level (students, families and all school personnel). Strategies for promoting whole-school mental health and wellness should include a social-emotional learning framework, mental health literacy, attention to stress management, the development of learning environments that meet the needs of all students and the establishment of positive social connections.

To learn more, visit the
 School Mental Health Resource and Training Center at
www.mentalhealthEDnys.org or call **518.434.0439**





MENTAL HEALTH EDUCATION LITERACY IN SCHOOLS LINKING TO A CONTINUUM OF WELL-BEING

New York State Education Department (NYSED) Amendments to Commissioner's Regulation (CR) Part 135 regarding Health Education

May 2018 the NYSED Board of Regents Permanently Adopted Proposed Amendments to Sections 135.1 and 135.3 of the Commissioner's Regulations; Education Law §804 was amended by Chapter 390 (Laws 2016) and Chapter 1 (Laws 2017) clarifying that a satisfactory program in health education in accordance with the needs of pupils in all grades that includes the several dimensions of health, is now required to:

- include mental health and the relation of physical and mental health; and
- designed to enhance student understanding, attitudes and behaviors that promote health, well-being and human dignity.

Schools may already be providing "mental health" education as previously stated in CR's Part 135, however this new legislation formalizes these new requirements in law.

To view the amendments, the NYS Mental Health Education Advisory Council's Panel Presentation to the Board of Regents, the Advisory Council Membership and the twenty-six recommendations please click on the following [NYSED Board of Regents Mental Health Education May 2018](#)

Mental Health Facts

Mental health is a critical part of overall health and *well-being* and is important throughout the life cycle; affecting thinking and learning, feelings and actions, influencing healthy decision making.¹

According to the World Health Organization "there is no health without mental health" with health being a state of complete physical, mental and social *well-being*; not only absence of disease.²

Education on the importance of the mind-body connection,³ and the several dimensions of health including mental health and the relation of physical and mental health, will enhance student understanding, attitudes and behaviors that promote health, well-being and human dignity.

"In childhood and throughout adolescence, mental health means attaining developmental and emotional milestones, learning healthy social skills and coping with challenging situations. Mentally healthy children/youth have a positive quality of life and function well at home, in school, and in their communities"⁴

According to the Centers for Disease Control and Prevention (CDC), "focusing on establishing healthy behaviors during childhood is more effective than trying to change unhealthy behaviors during adulthood."⁵

NEW YORK STATE MENTAL HEALTH EDUCATION ADVISORY COUNCIL (ADVISORY COUNCIL)

In August of 2017, NYSED, the New York State Office of Mental Health (OMH), and the Mental Health Association in New York State, Inc. (MHANYS), established the Advisory Council of over 75 expert cross-disciplinary and cross-sector partners, to develop resources, and recommendations supporting mental health education in schools.

Mental Health classroom resources developed by the Advisory Council identify skills and functional knowledge students should acquire, and tools to address stigma and promote best educational practices. These resources include an instructional guide, aligned with New York State (NYS) Learning Standards for health education in the classroom. The guide is reflective of skills and functional knowledge, scope and sequence for all levels (K-12) focused on health-related outcomes; and includes an instructional resource section.

WHAT SCHOOLS CAN DO

Review and assess current K – 12 health education curricula for alignment to new mental health education requirements;

Build capacity and strengthen relationships between educators and pupil personnel services (school psychologist, social worker, counselor, nurse);

Develop school-community partnerships with mental health professionals and organizations;

Identify strategies to engage families and students in supporting mental health and well-being;

Support a school climate “Culture of Care”; and

Leverage partnerships and build upon existing resources to develop a sustainable infrastructure for mental health.

Updated September 2018



NYSED'S NEXT STEPS

Continue a multi-prong, transparent, timely approach to address next steps.

Be supportive of, promote, and engage the [School Mental Health Resource and Training Center](#) developed by the Mental Health Association in New York State, Inc., to assist schools implement mental health instruction as part of the health education program.

December of 2018 - Advisory Council Meeting for a debriefing opportunity, and to discuss any lessons learned to improve ongoing implementation.

ADDITIONAL INFORMATION

NYSED's [Office of Curriculum and Instruction](#):
<http://www.p12.nysed.gov/ciai/health/>

[NYS Center for School Health](#):
<http://www.schoolhealthny.com>

NYSED's [Office of Student Support Services Social Emotional Learning](#): <http://www.p12.nysed.gov/sss/sel>

[Mental Health Association in NYS, Inc.:](#)
<https://mhanys.org/>
<https://www.mentalhealthednys.org>

NYS [Office of Mental Health](#):
<https://www.omh.ny.gov/omhweb/about/>

NYS [Office of Alcoholism and Substance Abuse Services](#):
<https://www.oasas.ny.gov/>

¹ <https://www.mentalhealth.gov/basics/what-is-mental-health>

² http://www.who.int/features/factfiles/mental_health/en/

³ <https://www.mentalhealth.org.uk/a-to-z/p/physical-health-and-mental-health>

⁴ <https://www.cdc.gov/childrensmentalhealth/basics.html>

⁵ <https://www.cdc.gov/healthyschools/wsc/index.htm>

A GUIDE FOR SUICIDE PREVENTION IN NEW YORK SCHOOLS



Table of Contents

ACKNOWLEDGMENTS	1
INTRODUCTION	2
UNDERSTANDING SUICIDE	3
Risk and Protective Factors	4
Warning Signs	6
HOW IS THE PREVENTION OF SUICIDE FRAMED?	7
TIER I UNIVERSAL INTERVENTIONS	8
Faculty and Staff Awareness Training	8
Focus on Social Emotional Competency Development (SEL)	8
School Connectedness/ School Climate	9
Upstream Prevention Programs	10
Mental Health Literacy Education	10
Student Suicide Prevention Education	11
Parent Education	11
TIER II TARGETED INTERVENTIONS	12
Groups at Higher Risk	12
Students Experiencing Stressful Life Events	14
TIER III INDIVIDUALIZED INTERVENTIONS	16
Determining Level of Suicide Risk	16
Assessment and Triage	16
Brief Safety Planning Intervention	17
Working with Parents	17
Working with Community Mental Health	18
Plan for Safe Return to School	18
In-School Suicide Attempts	18
Out-of-School Suicide Attempts	19
POSTVENTION	20
RESOURCES	21
REFERENCES	25

Acknowledgements

This guide was developed by the Schools and Youth Workgroup of the Suicide Prevention Council at New York State Office of Mental Health to provide best practice guidance and resources to schools toward the prevention of youth suicide. Written by Pat Breux and Mitchell Samet, the guide would not have been possible without the generous contributions of thought, expertise, and time of many people for which we are very grateful.

Schools and Youth Workgroup

Pat Breux

*Director of School & Youth Initiatives
Suicide Prevention Center of New York*

Sam Brinton

*Head of Advocacy and Government Affairs
The Trevor Project*

Rosa Gil

*President and CEO
Comunilife, Inc.*

Madelyn Gould

*Professor of Epidemiology in Psychiatry
Columbia University Medical Center*

Brianna Gower

*Director of YOUTH POWER!
Families Together in NYS, Inc.*

Glenn Liebman

*Chief Executive Officer
Mental Health Association in New York State*

Renee Rider

*Executive Director
New York State Council on Children and Families*

Mitchell Samet

*Workgroup Chair
New York Association of School Psychologist*

Victor Schwartz

*Chief Medical Officer
The Jed Foundation*

Peter Wyman

*Professor of Psychology
University of Rochester School of Medicine*

Additional Contributors and Reviewers

Scott Bloom, NYC Office of School Health

Rosa Cifre, Comunilife

Nancy Craig, Families Together in NYS

Peter Faustino, NYASP

Laura Ficarra, NYS OASAS

Sara Gorman, Jed Foundation

Brett Harris, NYS OMH

John Kelly, NYASP

Kathleen Liedka, NYS OMH

Sarah Miles, NYC Office of School Health

Keygan Miller, The Trevor Project

Amy Molloy, MHANYS, Inc.

Casey Pick, The Trevor Project

Kathryn Provencher, NYS OMH

New York State Education Department Suicide Prevention Workgroup

Maribeth Barney, NYSED

Donna Bradbury, NYS OMH

City of Albany School District Pupil Services Team

Kathleen DeCataldo, NYSED

Carrie Manchester, NYSED

Martha Morrissey, NYSED

Introduction

Recent media coverage of suicide deaths among high profile individuals has served to underscore the disturbing reality that suicide is a growing public health concern that does not discriminate on the basis of demographics. In fact, suicide has become the tenth leading cause of death in the US and has recently surpassed the death rate by motor vehicle accidents, homicides and breast cancer (NYSOMH, 2016). In 2017, over 47,000 individuals died by suicide in the United States, including 1,696 in NY State (CDC, 2019). New York State has one of the lowest suicide rates in the nation, at 8.6 per 100,000 as compared to the 13.4 per 100,000 national rate. However, given the high population, New York is among the top six states for the number of suicide deaths (CDC, 2019). Moreover, deaths by suicide in NYS have increased by 32% in the past decade, in stark contrast to gradual reductions in the death rate for other diseases such as cancer, heart disease or stroke (NYSDOH, 2019).

Each year, 4,600 children and teens, age 10-19, die by suicide. The rate of suicide in all age groups has continued to steadily rise in the last decade and the rate of suicide death among children 10 to 14 has doubled in that same time frame. As a result, in 2016, suicide was the second leading cause of death among young people ages 10-19 (CDC, 2018). According to the Centers for Disease Control and Prevention (CDC), among high school students in New York State in 2017, 17.4% seriously considered suicide, and 10.1% have made non-fatal suicide attempts (CDC, 2018).

Addressing the problem of youth suicide requires collaborative action across a variety of community agencies, but schools have logically assumed more of a leadership role in identifying, referring, and aiding youth with mental health needs. Schools also play a critical role in promoting psychosocial competencies that reduce vulnerability to suicide. The importance of schools is best highlighted by the recent adoption

of NYSED's 2018 Mental Health Education Literacy in Schools: Linking to a Continuum of Well-Being Comprehensive Guide. Given that the developmental trajectory for suicide risk can begin early in life, schools are uniquely positioned for building resilience among their students and developing a caring community within a positive school climate and culture necessary for the prevention of suicide. Since protecting the health and well-being of students is a fundamental role of all school staff, it is critically important that school districts develop policies and put procedures in place to reduce and respond effectively to suicide risk.

Given this critical responsibility, the following guidance document outlines best practices for school districts in New York State (NYS), to help protect the health and safety of all students and to guide school districts in developing policies and procedures to prevent, assess the risk of, intervene, and respond to youth suicidal behavior. This document utilizes the Multi-Tiered System of Support (**MTSS**) model, and includes a focus on universal supports, prevention and a positive school climate (Tier I), targeted supports for at risk groups and individuals (Tier II), and specialized supports and procedures for students in crisis or high risk (Tier III).

As related to suicide prevention efforts, effectively implementing the MTSS model includes three critical components for school districts: (1) Developing and implementing policies that includes suicide prevention, intervention and postvention. (2) Providing ongoing professional development for teachers and school staff on warning signs, risk factors, protective factors, and school prevention, intervention and postvention procedures. (3) Implementing educational programming and curriculum that integrates developmentally appropriate, social/emotional, suicide prevention, and mental health education as a basic foundation for health, safety and wellness for all students.

Understanding Suicide

The CDC defines suicide as *“Death caused by self-directed injurious behavior with an intent to die as a result of the behavior.”*

The operative word here is “intent”. If someone does something that is risky and dangerous, like drink and drive or take drugs and they die, it will be ruled an accidental death unless there is some evidence that they intended to die as a result of what they did.

There are many situations such as single car accidents with fatalities where there is a suspicion of suicide, but because the intent to die was never communicated, it will be ruled accidental. This also can help us understand why behaviors like cutting are called non-suicidal self-injury. Most of the time the self-injury is done with the intention of relieving some type of psychological pain - not dying. There is concern, however, that people who cut on multiple occasions lower their threshold for lethal self-injury. In other words, the more they get used to injuring themselves, the easier it might be to fatally injure themselves.

Another definition is sometimes helpful: “Suicide is an attempt to solve a problem of intense emotional pain with impaired problem-solving skills.” This definition captures what is happening to someone who is suicidal: they are experiencing a problem or a series of problems which are causing great psychological pain.

For example, if you think about a time in your life when you felt overwhelmed by what you considered to be a terrible problem, you can probably remember that your problem-solving was not at its best. Most suicidal people, including youth, do initially try to figure out how to cope with their problems. When that initial solution does not work and new problems get piled on, their coping skills start to diminish. Usually there is some type of triggering event that pushes them over the edge.

Understanding suicide from this behavioral perspective can help us appreciate that for most people, the decision to take their life is at the end of a process that has been going on for a while. While the final act of suicide may be impulsive, the thought process leading up to it is not. There is time to intervene and perhaps prevent the suicide if we know what to look for and how important it is to get them help.

DEFINITION OF TERMS

Suicide Ideation

Thoughts of ending one’s own life, regardless of how intense these suicidal thoughts are.

Suicidal Behavior

Any behavior resulting in an attempt or preparation for an attempt; this may include practicing or rehearsing for the attempt.

Suicide Attempt

Non-fatal self-directed potentially injurious behavior with any intent to die as a result of the behavior. A suicide attempt may or may not result in injury.

Non-suicidal Self-injury

Deliberate direct destruction or alteration of body tissue without a conscious suicidal intent (Erbacher, et al., 2015).

Postvention

interventions to address the care of bereaved survivors, caregivers, and health care providers; destigmatize the tragedy of suicide and assist with the recovery process; and serve as a secondary prevention effort to minimize the risk of subsequent suicides due to complicated grief, contagion, or unresolved trauma (Underwood, 2018).

Suicide Contagion

The phenomenon by which suicide and suicidal behavior is increased for some who are exposed to the suicide of others (HHS, 2019).


Risk and Protective Factors

Risk and protective factors for suicide can be envisioned through the lens of resiliency theory (Zimmerman, 2013), which tells us that suicide is complex behavior that is influenced by the dynamic interplay of multiple risk and protective factors. These risk and protective factors can be understood as occurring along a social ecology continuum that includes individual, relationship and cultural/environmental factors (CDC, 2002).

This continuum starts at the **individual** level, encompassing a person's biology, psychology, and social history. Risk factors on this level include depression, impulsive or aggressive responses to stress, poor coping skills, substance use, problem gambling, and prior experiences with death and trauma.

Risk is also influenced by a person's **relationships** with family, peers, and mentors. These relationship factors include experiences such as interpersonal violence, abuse or sexual assault, bullying, family separations, the loss of an important friend or family member, no access and/or lack of engagement with mentors or having those close to you not recognize or be aware of your help-seeking attempts/signals.

On the continuum are **cultural/environmental** factors, including the perception of social acceptability of suicide in popular culture or among a peer group, codes of silence that discourage reaching out to adults for assistance, having access to the means to kill oneself, poor coordination of services, limited access to care, or a lack of social connectedness. The work of prevention is to identify and implement interventions that decrease risk factors or increase protective factors at varying points along the social ecology.



Concentric Circles of the Social Ecology Continuum (WHO,2014)

RISK FACTORS

Individual

- Mental disorder, especially depression
- Previous suicide attempt
- Alcohol & substance use and misuse
- Feelings of hopelessness
- Belief that suicide is a viable option
- Impulsive or aggressive reactions to stress
- Chronic or intense stress

Relationship

- Physical, psychological, sexual abuse
- Physical or sexual assault
- No mentors
- Loss of important relationship
- Rejection by peer group
- Loss of family member or friend to suicide
- Peer group engaged in self-destructive behavior
- Peer group endorses suicide as acceptable
- Highly stressed family
- Bullying
- Help-seeking signals ignored

Environment

- Suicide contagion
- Social rejection
- Community violence
- Poor coordination of community resources
- Access to means of suicide
- Access to environments where others have died
- Mental health care is stigmatized

PROTECTIVE FACTORS

Individual

- Good emotional self-regulation
- Easy-going personality
- Social problem-solving skills
- Positive activities
- School success

Relationship

- Access to mentors
- Family support (Parent involvement)
- Peer group engaged in pro-social behavior
- Caring friends
- Supportive relationships with mental health and medical providers

Environment

- Effective bully prevention practices
- School connectedness
- Effective care for mental and substance use disorders
- Opportunities to experience success, generosity
- Help-seeking is encouraged
- Coordinated community services
- Access to mental health and health care

(Breux, 2017)

Warning Signs

It is important to remember that most risk factors are common, and in and of themselves are no more predictive of a suicide than risk factors for heart disease predict heart attacks. Risk factors are simply predictors of potential risk, but a percentage of students exhibiting risk factors may start to exhibit suicide warning signs.

Stressful life events or situations can serve as triggering events for suicidal behavior. For young people, these precipitants may include events like getting in a fight with a friend or family member,

getting in trouble at school, a bullying incident, family separation, illness or death in the family, or loss of a friend or opportunity. Real or perceived losses, and upcoming events for which a student feels unprepared and overwhelmed, can also be precipitating events. When stressful life events are coupled with multiple or significant risk factors, it is important that school staff can recognize warning signs for suicide. Warning signs are observable signals that tell us to stop and pay immediate attention to a particular student because they may be at risk for imminent suicidal behavior. The word **FACTS** serves as a helpful acronym for identifying the most commonly recognized warning signs of suicide:

F is for FEELINGS. Hopelessness, worthlessness, despair about the future, or excessive worry.

A denotes ACTIONS. Actions include trying to gain access to a gun or pills, behaving recklessly or increasing alcohol or drug use. Actions can also include showing aggressive behavior that is inconsistent with the student's previous demeanor, self-harming behaviors, or being involved in bullying or looking online for ways to die.

C indicates CHANGES. This is a very important category because it means we are looking for changes from the student's previous attitude, moods, or behaviors which have been noticeable for at least a couple of weeks. For example, students who were active may become withdrawn, quit athletic teams, stop paying attention to personal appearance, daydream or even fall asleep in the classroom. It would be impossible to list all the potential behaviors you might see, so simply concentrating on recognition of changes from previous behaviors is a real key to making assessments in this category.

T represents the THREATS that some students make or hint at. These can be specific statements of intent like "I'm done with living" or "I'm thinking of killing myself" or worrisome innuendos in writing or other class assignments like "People would be better off without me". Threats may also be posted on social media sites. Whether specific or vague, threats tell us the student is thinking about death or suicide, and this escalates the level of concern.

S refers to SITUATIONS that may serve as triggers for the suicide. These include getting into trouble at home, in school, or with legal authorities; personal losses in relationships, opportunities, or even losses of less tangible things like self-esteem or hopes for the future. Life changes for which the student feels overwhelmed or unprepared such as moving or the transition after high school graduation can also serve as triggers. The most worrisome time is between the occurrence of the triggering situation and its resolution—in that period of uncertainty before the outcome is known (SPC-NY, 2019).


**This version of FACTS is taken from Suicide Safety for Teachers and School Staff, Lifelines Prevention, and Act on FACTS: Making Educators Partners in Youth Suicide Prevention.*

How is the prevention of suicide framed?

School districts play a central public health role in combating infectious diseases, malnutrition, community violence, accidental injuries, substance use and/or problem gambling addictions, and sexually transmitted infections by implementing such measures as health screenings, school nutrition programs, and health and safety education. These efforts seek to protect school communities through health education, prevention screenings, and intervention across an entire school's population. In each of these cases, schools have utilized a multi-tiered approach.

As the risk factors for suicide are numerous and their relationship with suicide risk is complex, the prevention of suicide requires that multiple strategies be employed. **The World Health Organization** calls for the use of a multi-tiered approach when addressing suicide prevention. The MTSS model is familiar to most school professionals, given that it is utilized for many school-based prevention initiatives, including academic concerns, bullying prevention, school climate, or mental health education literacy.

- **Tier I** interventions include universal-level interventions. At this level, we are looking proactively at the awareness and training needs of all members of our school community including parents, assessing our school environment and school climate for how protective factors can be enhanced, and considering upstream prevention programs. Tier I interventions alone are appropriate for roughly 80% of the population.
- **Tier II** interventions include selective strategies targeting 1) groups who may be at higher risk, 2) students exhibiting warning signs, and 3) students experiencing stressful life events that may put them at elevated risk. Interventions at this level will include social supports and key target points in time when screening for suicide risk may be prudent. Tier II interventions combined with Tier I supports are appropriate for 15% of the population.
- **Tier III** interventions are focused on individual students that are acutely affected by a suicide loss, are engaging in suicidal behavior, or are demonstrating acute suicide risk. Tier III interventions combined with Tier I and Tier II supports and combined, are appropriate for the remaining 5% of the population (Schaffer, 2017).



Tier III - Individualized Supports

- Determine level of risk
- Notify Parents
- Make a safety plan
- Refer to community mental health
- Follow up, return to school meeting

Tier III - Target Group Supports

- Screening, interventions, and supports for higher risk groups
- Screenings and supports during stressful life situations

Tier III - Universal Supports

- Faculty and staff awareness education
- Focus on social emotional learning
- School connectedness
- Mental health literacy education
- Student suicide awareness education

The Suicide Prevention Center of New York (SPC-NY) offers a free **School Readiness Workbook** to assist in the suicide prevention planning process. Additionally, the **Comprehensive Developmental School Counseling/Guidance Plan** can support evidence-informed suicide prevention practices. Suicide prevention programming can be considered in the development of this district-wide planning process which is designed to operationalize comprehensive multi-tiered programming to support the emotional and behavior well-being of young people.

TIER I Universal Interventions

School districts should develop policies and programs which include Tier I universal suicide prevention. One way to envision suicide prevention in school is through the lens of a “**Competent and Caring School Community**” (Underwood, 2010). This setting is one in which all members of the school community play a role in prevention, from the school board to parents, staff, and to students. Everyone is responsible for the safety and well-being of each other and have the competency to recognize when to ask for help and know how and where to access assistance if there is a concern about suicide.

Faculty and Staff Awareness Training

It is critical that the district develop written procedures for the staff to follow, when warning signs of suicide are observed or suspected. Staff awareness training should include how to identify warning signs and the procedures to follow when referring a student thought to be at risk for suicide. Faculty education must also include guidelines about what educators can do to promote protection from suicide including the promotion of positive mental health, how to handle suicide related topics in the classroom, and how to identify developmentally appropriate materials for classroom use and school/community resources.

High quality resources for staff awareness training are readily available. The Society for the Prevention of Teen Suicide provides a free online two-hour course, *Making Educators Partners in Youth Suicide Prevention: Act on FACTS*. A **three-minute trailer** is available and the training modules can be accessed by registering at **SPTS University**. *Suicide Safety for School Staff*

is a brief training (60 to 90 minutes) designed by the Suicide Prevention Center of New York (SPC-NY) to meet the basic awareness needs of faculty and school staff. A network of over 200 trainers are available and connected to 10 regional training centers across New York. To find a trainer or to learn more about this initiative, contact **SPC-NY**. New York City educators also have the option of accessing free online modules of **Kognito’s At Risk**.

Focus on Social Emotional Competency Development (SEL)

Extensive research indicates that effective mastery of social emotional competencies is associated with greater well-being and better school performance. On the other hand, the lack of competency in these areas can lead to a variety of personal, social, and academic difficulties. In fact, a study of young students found a significant relationship between students’ social emotional competencies in first grade and their outcomes 13 years later (Bradshaw, 2013). Those students with early pro-social skills were more likely to graduate from high school on time, complete a college degree, and achieve and maintain full-time employment. Further, during high school they were less likely to be involved with police, use alcohol or misuse opioids, or require medication for emotional or behavioral issues. In addition to the many other benefits of SEL, lifetime suicide risk is diminished when students learn social problem-solving skills and learn to cope with emotional challenges and life stressors. Effective SEL programs have demonstrated an increase in social connectedness, which is an additional protective factor against suicide risk.

In 2018 and 2019, the New York State Education Department (NYSED) published new resources to assist schools and districts in implementing SEL strategies through explicit instruction, integration with academic curricula, general teaching practices, integration of SEL in school and district policy and practice, and professional development and SEL support for adults in the school community. These resources can be found on NYSED's SEL website, and include: (1) SEL benchmarks for voluntary implementation, (2) a framework document explaining SEL concepts and the need for and benefit of SEL, (3) a guidance document to provide implementation strategies and resources for districts and schools; and (4) district-developed crosswalks aligning SEL competencies, academic standards, classroom activities, and general teaching practices. **Social Emotional Learning: Essential for Learning, Essential for Life** provides a framework explaining SEL concepts, and the need for and benefit of SEL to assist schools with implementation of the NYSED recommended SEL benchmarks. **The Collaborative for Academic, Social, and Emotional Learning (CASEL)** provides an extensive collection of resources, including a **Guide to Schoolwide Social and Emotional Learning**, to assist schools and districts in the implementation of Social Emotional Learning strategies.

School Connectedness/ School Climate

School connectedness is best defined as a belief held by students that adults and peers at school care about them and about their learning. School connectedness leads to both positive educational and health outcomes which includes a sense of student belonging and contribution, believed to be important protective factors against suicide risk. CDC developed **School Connectedness: Strategies for Increasing Protective Factors Among Youth**, a resource document for schools.

School climate is defined as the way school culture affects a child's sense of safety and acceptance, and consequently is a critical determinant of their ability to focus on the task of learning. Measuring school climate is a crucial step in creating a safe and caring environment. Collecting data through surveys of students, parents, and staff is an important component of a school climate assessment. The NYSED promotes efforts to build and maintain positive and healthy school climates, and has engaged in a **School Climate Pilot Project** using United States Department of Education School Climate Surveys.

Upstream Prevention Programs

Research suggests that universal prevention programs for children may effectively reduce adolescent emotional and behavioral problems that are risk factors for future suicidal behavior. While many upstream prevention programs, such as Positive Behavioral Interventions and Supports (PBIS) framework, Second Step program and mindfulness training, may enhance protection against suicide for students, few programs have actually been studied for this outcome. Two programs, the **PAX Good Behavior Game**, and **Youth Awareness of Mental Health** are universal interventions that have shown to reduce suicide attempts over time. **Sources of Strength** is a program that utilizes trained student peer leaders and adults and has been shown to improve school-wide help-seeking and engagement of adult help for suicidal peers, with the greatest benefits for suicidal youth (Wyman, 2010).

Mental Health Literacy Education

Mental health literacy is defined as “knowledge and beliefs about mental disorders which aid their recognition, management, or prevention.” A growing body of research in mental health literacy suggests that an increased understanding of mental health will improve help-seeking behaviors, reduce stigma, and promote better health outcomes (Kutcher, 2016). This is true for both young people and their supporters such as families, teachers, and peers.

Health education that respects the importance of mental health, as well as the challenges of mental illness, will help young people and their families and communities feel more comfortable seeking help, improve academic performance and, most importantly, save lives. According to the CDC, “focusing on establishing healthy behaviors during childhood is more effective than trying to change unhealthy behaviors during adulthood.” An equally important part of this conversation is to help students identify risk and protective factors, as learning and resiliency can result in positive decision-making and life-long success, which are the primary goals of health and education. In June, 2018, NYSED published a comprehensive

guide **Mental Health Education Literacy in School: Linking to a Continuum of Well-being** to provide guidance for developing effective mental health education instruction in the classroom. The guidelines comply with the recent amendments to Education Law 804, while also looking at embedding mental health well-being within the broader context of the school environment to promote a positive culture and climate.

The School Mental Health Resource and Training Center, was established by the Mental Health Association in New York State, Inc. (MHANYS) to help schools comply with the new law. The Center provides free online mental health training (CTLE-Eligible) and instructional resources for educators, as well as mental health resources for parents and guardians, students and community-based mental health providers. Statewide, Resource Center staff can provide professional development for school staff, including MHANYS Mental Health and Wellness 101, mental health instruction for students, and technical assistance for developing mental health curriculum and schoolwide mental health promotion strategies. Finally, the Resource Center can provide support in establishing school-community partnerships.

MHANYS addresses suicide prevention in their white paper **Mental Health Education in New York Schools: A Review of Legislative History, Intent, and Vision for Implementation**. This document offers core elements that should be included in school mental health curricula and acknowledges that unrecognized, untreated and late treated mental illness elevates the risk of mental health disabilities, suicide, and self-injury (Richter, 2017).

Additionally, there are several New York State community partners prepared to assist schools in promoting mental health literacy including the **JED Foundation** which licensed a **mental health curriculum**. In addition, **teenmentalhealth.org** is an evidence-based, comprehensive mental health curriculum for grades 8-10 that is implemented through a train-the-trainer model. The training has been shown to significantly increase educators' knowledge about

mental health and significantly decrease stigma. National Alliance on Mental Illness NYS (**NAMI NYS**) offers **Ending the Silence** presentations for students, educators and families, a stigma reducing program. The American Foundation for Suicide Prevention (**AFSP**), which has chapters across NYS, offers **More than Sad** for students, educators and families, a video-based educational presentation on teen depression.

Student Suicide Prevention Education

Suicide prevention education can begin in middle school or high school. It should include common warning signs and how and when to seek help. However, it should be noted that not all suicide prevention education efforts are helpful. Presentations can inadvertently stigmatize a student who may need help. In some cases, sensational presentations by persons with a compelling story have raised concerns about suicide contagion and about students learning how to kill themselves from such presentations. Yet, suicide prevention education can improve student knowledge of how, when, and where to seek help. When implementing suicide prevention education, it is important that the educator be well trained in evidence-based suicide prevention that is developmentally appropriate, that the presentation be integrated with other similar content rather than standing alone, and that it be done in a classroom or small group, rather than in an auditorium presentation. Classroom teachers are wise to collaborate with student support service providers in the development of the presentation. When student support service providers co-teach or support classroom teachers, they can assure that students know how to access help. Since the best outcome of any such education would be that students would recognize signs of distress in themselves and their classmates and reach out for help, it is best that parents and adults in school have also had suicide prevention awareness education.

The SPC-NY offers a **tip sheet** for classroom presentations. Several programs and curriculum exist for student suicide prevention awareness. As an example, **Lifelines Prevention** is a manualized curriculum from Hazelden Publishing that has lesson plans for middle and high school students. Another program, **Sources of Strength (SOS)** engages peer

leaders to enhance protective factors across a high school population. SOS has been shown to improve social norms associated with suicide prevention with the greatest changes being among students with suicidal ideation. Major findings include improved perceptions of support from adults and increased likelihood that they would tell an adult if they were concerned about suicide (Wyman, 2010).

Parent Education

Parents need information about the prevalence of youth suicide, its warning signs, how to respond if their child or another child is at risk, and where to turn in their community for help. Some parents do not relate well to written materials or videos, especially on difficult topics like suicide and may respond better to verbal or personal communication.

The Society for the Prevention of Teen Suicide offers several resources that can be helpful for designing parent awareness or education activities. Among their resources is a free 17-minute video titled **Not My Kid** designed to answer the questions most frequently asked by parents about suicide. The Mayo Clinic offers a 5 minute **video** that focuses on the warning signs and how to respond. The Suicide Prevention Center of New York offers a **parent brochure** that can be downloaded or linked.

TIER I CONSIDERATIONS

- Written procedures should clearly delineate how to refer a student when suicide risk is suspected and should be reviewed annually with all staff.
- All school staff should have basic suicide awareness training and annual refresher training on warning signs and how to respond.
- School climate surveys of students, parents, and staff should be performed regularly, and the data should be used to inform school improvement planning.

TIER II Targeted Interventions

As noted above, Tier II interventions include selective strategies targeting (1) groups who may be at higher risk, and (2) those experiencing stressful life events that may put them at elevated risk. Below, are some, but not all groups that research has demonstrated have a higher risk of suicide (King, 2013). In discussing groups that may be at higher risk, it is important to question what it is about being in that group, that might lead to increased risk. It is these circumstances for which school health and mental health professionals including school psychologists, social workers, school counselors, nurses, or on-site community mental health providers can then consider appropriate interventions.

While the characteristics that define a student as a member of a higher risk group might be a fixed fact and may or may not be alterable, a focus on stressful life events provides a slightly different lens. If we recognize that the anticipation and aftermath of a stressful life event, particularly those that represent a meaningful loss to the individual, can be a critical time for suicide risk, we can preemptively intervene with supports and consider opportunities for screening and recognizing warning signs of suicide.

Groups at higher risk

- LGBTQ+ (Robinson, 2012)
- Latina youth (Zayas, 2009)
- Students with depression (AFSP, 2019)
- Students using drugs or alcohol (AFSP, 2019)
- Students with a previous attempt
- Students with emotional dysregulation
- Student with a history of self-injury (Klonski, 2014)
- American Indian or Alaska Native
- Students with intellectual disabilities (Ludi, 2013)
- Students with health problems and physical disabilities (Jones, 2008)

Stressful life events that may increase risk (AFSP, 2019)

- Family separation
- Death of a friend or family member
- Death of a parent
- Exposure to violence
- Disciplinary action
- Failure
- Loss of home
- Exposure to the suicide of another
- Engaged in bullying

Groups at Higher Risk

Lesbian, Gay, Bisexual, Transgender, Queer/ Questioning (LGBTQ) Youth

Students identifying as lesbian, gay or bisexual may be 4 to 5 times more vulnerable to suicide risk than their heterosexual peers. Lesbian, gay or bisexual youth are also more likely to experience discrimination, bullying, violence, and victimization. However, victimization alone does not explain the elevated risk (Robinson, 2012). The risk for attempts in this group is significantly lessened if the student is in a supportive social environment (Hatzenbuehler, 2011). A student who feels they can talk about their problems to just one school staff member is 30% less likely to report making multiple suicide attempts than youth who do not have a safe adult to talk to, highlighting the importance of training school staff to support LGBTQ youth with sensitivity and cultural competence (AFSP, 2019).

Transgender adolescents and gender non-conforming adolescents as a demographic group experience high rates of suicide attempts (Becerra-Culqui, 2018). It is recommended that school personnel be educated about the risk and that social supports are critically important. Because harassment, discrimination, and victimization are implicated in elevated suicide risk for this group, it is critical that schools have policies in place that create safe and supportive school environments. In July of 2015, NYSED published **Guidance to School Districts for Creating a Safe and Supportive School Environment for Transgender, and Gender Nonconforming Students**.

The Trevor Project is a national resource that offers online learning modules as a teacher resource for middle schools and high school classrooms. They also offer 24-hour crisis support via text, chat, and phone for LGBTQ youth experiencing emotional distress.

Latina Youth

Latina adolescents report feeling sad and hopeless at higher rates than their black or white classmates. They also have higher rates of serious suicide attempts requiring medical attention (CDC, 2018).

Comunilife's Life is Precious (LIP) program provides culturally and linguistically appropriate services for Latina teens, who are living with depression and/or have seriously considered or attempted suicide and their families. LIP provides educational support (tutoring and school system navigation), creative art therapies, wellness activities, case management and concrete family services. Preliminary evaluation demonstrated LIP is effective in reducing suicidal ideation and depression (Humensky, 2016).

Students with Emotional Dysregulation and Self-Injurious Behavior

The relationship between non-suicidal self-injury (NSSI) and suicide is complex. Research demonstrates that non-suicidal self-injury is an important risk factor for a suicide attempt (Klonsky, 2014). Additionally, a previous suicide attempt is an important risk factor for death by suicide. Both risk factors can be viewed as indications of emotional pain and increased capability to engage in self-harm that may lead to death.

Emotional dysregulation related to a chronic or situational stressor, or a child's vulnerability to distress can lead to maladaptive behavior patterns in certain vulnerable children. Essential life skills for such students include: 1) Awareness, understanding, and acceptance of emotions, 2) Ability to engage in goal-directed behaviors and inhibit impulsive behaviors when experiencing negative emotions, 3) Use of situationally appropriate strategies to modulate the intensity or duration of emotional responses rather than eliminating emotions entirely and 4) Willingness to experience negative emotions as part of pursuing meaningful activities.

DBT in Schools is a promising program. It is an adaptation of an effective therapeutic clinical intervention Dialectical Behavior Therapy (**DBT**) that has been proven to decrease suicidal behavior and suicidal ideation. It includes lesson plans for classroom social/emotional skills development (Dadd, 2016).

Youth with Mental Health and/or Substance Use Disorders (SUD)

Mental health conditions: depression, attention-deficit hyperactivity disorder, eating disorders, intermittent explosive disorder, and conduct disorder are important risk factors for suicidal behavior among young people. An estimated one in five children have a diagnosable mental disorder that can result in significant impairment; therefore, school staff may play a pivotal role in recognizing and referring the student to treatment that may reduce risk and improve longer-term outcomes.

Suicide is the leading cause of death among those with substance use disorders and 20% of suicide deaths involve alcohol. Acute alcohol intoxication is present in 30 to 40% of all suicide attempts. The disinhibition that is associated with alcohol consumption is one factor. When a student with suicide risk is also using alcohol, there should be increased concern for the possibility of a suicide attempt (SAMHSA, 2016).

Screening, Brief Intervention, Referral to Treatment (SBIRT) is a framework that takes a comprehensive approach to address substance use. Many local prevention providers across New York can partner with schools to offer this evidence-based practice. The NYS Office of Alcohol and Substance Abuse Services (OASAS) offers a Directory of local Substance Use Prevention Programs. Teen Intervene is a brief early intervention program for purchase for students ages 12- to 19-years who display mild to moderate problems associated with alcohol or drug use.

Students Experiencing Stressful Life Events

Youth in Out-of-Home Placement or Experiencing Homelessness

Youth involved in the juvenile justice or child welfare systems have a high prevalence of many risk factors for suicide. As much as 60 to 70% of young people involved in the juvenile justice system meet criteria for at least one psychiatric disorder, and youth in juvenile justice residential programs are three times more likely to die by suicide than the general youth population (Scott, 2015). According to a study released in 2018, nearly a quarter of youth in foster care had a positive diagnosis for major depression in the last year. Additionally, a quarter of foster care youth reported attempting suicide by the time they were 17.5 years old (Okpych, 2018).

For youth experiencing homelessness, rates of self-injury, suicidal ideation, and suicide attempts are over two times greater than those of the adolescent population in general. These young people also have higher rates of mood disorders, conduct disorder, and post-traumatic stress disorder. One study found that more than half of runaway and homeless youth experience suicidal ideation (Barnes, 2018).

Bullying and Suicide

There is ample empirical evidence that bullying behavior and suicide are closely related. Youth who report any involvement with bullying behavior are more likely to report higher levels of suicide related behavior, as compared to youth who do not report bullying (CDC, 2014). However, the relationship between bullying and suicide is complex. Persistent bullying can lead to or worsen feelings of isolation, rejection, exclusion and despair. While studies have shown that young people who are bullied and those who bully others are at heightened risk for suicidal behavior, youth who exhibit both pre-existing risk for suicide and who are concurrently involved in bullying or experiencing other negative life events are at highest risk.

Researchers at the University of Buffalo's **Alberti Center for Bullying Abuse Prevention** demonstrate that being a perpetrator of bullying is linked to depression, suicidal ideation, and suicide attempts (Espelage, 2013), as well as negative perceptions of school climate (Nickerson, 2014). The Center offers evidence-based tools to effectively change the language, attitude, and behavior of educators, parents, students, and society.

Educators can reduce suicide risk through improved bullying prevention initiatives, increased focus on school connectedness, and the use of evidence-based social/ emotional approaches as illustrated in Tier I above. It is imperative to convey safe and accurate messages about bullying and suicide to youth, especially to young people who may be at risk for suicide.

It is critical to understand that, suggesting that suicide is a natural response to bullying or providing repeated opportunities for at-risk students to see their own experiences of bullying, isolation, or exclusion reflected in stories of those who have died by suicide, can increase **contagion** risk by contributing to thoughts that frame suicide as a viable solution. Idealizing young people who die by suicide after being bullied, or creating an aura of celebrity around them, may contribute to an at-risk youth's illogical thoughts that suicide is the only way to have a voice or to make a difference for others. However, when school personnel know that a student is involved in bullying, they should not hesitate to ask students direct questions about hopelessness and thoughts of suicide. Whenever possible, discussions on bullying and suicide should center on prevention (not statistics), resiliency, and encourage help-seeking behavior (CDC, 2014).

Death of a Parent

Approximately 4% of children in western countries including the US experience the death of a parent during childhood, and parent death is rated as the most stressful event to happen to a child. Although most children adapt well (75-80%), parentally-bereaved children are at increased risk for problems into adulthood including risk for suicide (Guldin, 2015). The Family Bereavement Program which teaches coping and resilient parenting demonstrated long term reductions in suicidal ideation and attempts (Sandler, 2016).

Students Returning from Psychiatric Hospitalization

This is a particularly stressful time and for which special consideration to safety planning and supports should be given. It is discussed more fully in the Tier III section below.

TIER II CONSIDERATIONS

- Policy to delineate when a student may be at risk and need to be screened for suicide risk. Such times might include when a student is:
 - Exhibiting warning signs
 - Identified by staff to be at risk for suicide
 - A victim of a bullying incident
 - Receiving serious disciplinary action
 - Bereaved or exhibiting emotional distress
 - Being referred for mental health care
 - Self-injuring
- Social supports should be accessible to students and their families.
- When conducting climate surveys, special attention should be paid to groups that may feel marginalized or discriminated against and additional supports should be considered.

TIER III Individualized Interventions

Helping a student at risk for suicide is a collaborative process, one in which parents, community providers, school professionals, and the student all have a role to play and must work together to keep a student safe. Intervention should be undertaken with developmental competency and an attempt to understand cultural concerns. The process outlined below includes assessment and triage, brief safety planning intervention, collaborative decision-making and referral to community providers, and includes a plan for a student's safe return to school.

Tier III intervention is for students identified as exhibiting warning signs and for whom there is some identified concern for suicide. Standardized written procedures for intervening when a student is identified to be at risk for suicide is a considerable asset for those charged with assisting students who are struggling. However, given that suicidal behavior is hard to predict, and the safety concerns are so significant, it is critical that all school health, mental health, and designated administrative staff, have adequate and ongoing professional development and support to implement standardized procedures with fidelity.

As noted above in Tier I, teachers and other school staff play a critical role in recognizing warning signs in students and connecting at-risk students to the identified school-based health and mental health professionals who can assess the level of risk and intervene appropriately. Every school district should have a protocol in place for referring students who may be at risk.

Determining Level of Suicide Risk

Any student exhibiting any of the aforementioned FACTS warning signs, should receive an immediate screening for suicide risk by school-based health or mental health professionals who have received appropriate training. The six question **Columbia Suicide Severity Rating Scale (CSSRS)** is a recommended screening instrument. **Numerous studies** support the psychometric properties of the CSSRS, including: its sensitivity to change, internal consistency, inter-rater reliability, and cross-cultural and multilingual application. Research demonstrates its effectiveness as a tool for preventing suicides.

In developing suicide prevention policies, school leaders should identify who conducts such screening, what instrument to use, how it should be documented, how the information from the screening should be communicated, and who has access to it. Additionally, all staff need to be provided with training on the procedures to follow if a student is identified as exhibiting warning signs.

Assessment and Triage

As a first step, standardized procedures should be in place regarding the assessment and triage of students who may be at risk for suicide. Once a student is identified at risk for suicide, the student should *remain* under supervision until appropriate level of risk is assigned and intervention is in place. Assessment for suicide risk is best understood as happening in the midst of a semi-structured interview. Students often react better to a discussion of what is bothering them than to a rote clinical interview. Prepare to meet the student with calm reassurance and collect additional objective data ahead of time if possible, including grades, attendance, disciplinary referrals, and medications. Have as much information as possible about the reason for the referral. It can be helpful to directly address any resistance by the student. With younger children, they may need reassurance that they are not in trouble.

The goal of suicide risk assessment by school-based health and mental health professionals should be for the purpose of triage and safety planning. As part of the semi-structured interview, use of a brief, standardized tool with some evidence of reliability and validity is recommended. The CSSRS is one such tool. It is offered in both brief (six question) and long form versions and the language can be modified to be culturally or developmentally relevant. The tool is available in 114 languages and can incorporate information gleaned from other sources than just the interview. The tool and training on the use of the tool are offered free of charge and accessible online at the **Columbia Lighthouse Project**. Another advantage to this tool is that it can be administered by various professionals and it has been proven to be more accurate than a clinical interview. The six question CSSRS also helps with triage decisions. Typically, schools stratify risk into low, moderate, or high risk and designate a checklist of minimal interventions at each level.

Advantages to a Second Person

It is recommended, whenever possible, that intervention is achieved with two professionals. One who is looking out for the physical and emotional needs of the student and providing constant supervision, while the other is more focused on the logistical and communications aspects of the intervention.

Helping Students at Risk for Suicide

Helping Students at Risk for Suicide (HSAR) is a model for school-based suicide intervention that utilizes a collaborative approach. HSAR was developed through a partnership between the New York Association of School Psychologists (NYASP), SPC-NY, and the Alberti Center for Bullying Abuse Prevention at University of Buffalo. HSAR incorporates evidence-based tools including the CSSRS and the Stanley/Brown Suicide Safety Plan.

Brief Safety Planning Intervention

While a student benefits when school and community mental health professionals can collaborate in planning to assist a student, it is recognized that brief interventions at school can also be helpful.

Safety Planning Intervention is an evidence-based intervention in which a mental health professional works collaboratively with a student to think through coping strategies and resources that they can access to help keep themselves safe from suicide. Students benefit from having supports that are customized to their individual needs in the school environment (Stanley, 2012).

Additionally, students benefit when school professionals can competently discuss access to medications, weapons, and other means of self-harm with students and their parents. **CALM** training prepares professionals for this conversation. A means reduction discussion is an important part of a school safety planning meeting.

The next step in any intervention is to ensure that students and their parents have access to 24-hour crisis services. While every community in New York should have some access to emergency mental health services, it is important for school leaders to reach out and understand what the resources can and cannot do before referring families in crisis. Additionally, the

National Suicide Prevention Lifeline at 1-800-273-8255 and the **Crisis Text line** (Text "Got5" to 741-741) provide access to emotional support 24 hours a day. For residents of New York City, **NYCWELL** is an additional resource.

Working with Parents

Parents or guardians should be contacted as soon as possible after a student has been identified as being at risk for suicide. The person who contacts the family is typically the principal, school psychologist, social worker, school counselor, or a staff member with a special relationship with the student or family. Staff need to be sensitive toward the family's culture, including attitudes towards suicide, mental health, privacy, and help-seeking. It can be helpful to recognize that while the school knows a more public side of a student's life, parents know more about the student's personal life. It is through the combination of these two perspectives that the best understanding of the situation and consequently how to help can be reached. Garnering information from the family by going slow, demonstrating empathy and an interest in understanding a parent's perspective, and asking open ended questions can set the stage for collaboration. Questions should impart the school professional's desire to partner with the family on the collective goal of helping and keeping the child safe.

When meeting with parents or guardians:

1. Describe the situation and ask that they come to school immediately.
2. When the parent/guardian arrives, explain why their child is thought to be at risk for suicide.
3. Explain the importance of securing all firearms, medications, alcohol, and other dangerous items.
4. Explain options for follow up with mental health services or further emergency evaluation based on level of risk. Provide information about services and if possible assist with making appointments.
5. Ask parents/guardians to sign a form acknowledging that they were notified about concerns, recommendations and plan for follow up.
6. Identify the staff member who will be following up with parents/guardians and discuss when the follow-up will occur.
7. Explain that a return to school meeting will be scheduled for safety planning and support.
8. Request permission to share information with a health care or mental health care provider.

Family support is critical in ensuring a student is safe from suicide risk. And yet, families are often unprepared and without the professional supports they may need to deal with the complex emotions, shock, or distress they may encounter when notified that their child is at risk for suicide. Parents often need support and assistance to come to terms with their child's risk and their reaction to it. *Adapted from "Preventing Suicide: A Toolkit for High Schools" pg72 (SAMHSA,2012).

Families Together in New York State (FTNYS) is a state wide, parent-governed, non-profit organization that represents families of children with social, emotional, behavioral health and cross-systems challenges. FTNYS has chapters in all regions of New York State, and offers peer support, advocacy, and education for families navigating the mental health system.

Working with Community Mental Health

Students are safest when the mental health professionals at school partner with mental health professionals in their communities. While some schools benefit from strong relationships with community mental health providers, other schools struggle to find accessible mental health care for students. Much work is being done on the state and national levels to ensure that mental health services are more accessible to students. New York State Office of Mental Health offers guidance and resources for schools and mental health services to improve access to care, Improving School and Community Outcomes for Children and Adolescents with Emotional and Behavioral Challenges.

Systems of Care (SOC) initiatives are a growing resource in communities across New York State. These interagency committees, consisting of mental health and social service resource providers, meet to discuss the needs of children, youth, and families with a goal of improving access to care. They can be a great resource to schools as brainstorming partners to develop solutions to meet immediate or emerging needs. The Conference of Local Mental Hygiene Directors' website http://clmhd.org/contact_local_mental_hygiene_departments/ provides contact information for Single Point Of Access (SPOA) Coordinators and Local Mental Hygiene Directors, who in many cases are coordinating SOC initiatives. Contact them to inquire about the existence of a local SOC committee and how they may be a resource to your school community. More information on the SOC framework may be found on www.msnavigator.org.

Plan for Safe Return to School

When a student returns to school from a mental health referral in the community, after medical treatment for a suicide attempt, or after psychiatric hospitalization it is a crucial time for ensuring that appropriate supports are in place.

Ideally a return to school meeting is scheduled in which current suicide risk can be assessed, safety plan, current medications, and mental status can be reviewed. Students may need academic supports or accommodations if there has been a significant absence or if there are residual changes in cognitive reserve for learning related to new mental health diagnoses or injuries. Students may require assistance navigating the social milieu at school. Are there environments, classes where they might more likely become emotionally distressed and do they have strategies in place for managing this? They also need to have a schedule for check-ins with school-based mental health professionals and ways to access supports in a hurry when needed.

Students may need assistance sorting out what to say about their suicidality and to whom. They may need help managing rumors or explaining their absence or explaining visible injuries, which can also be a source of considerable distress. The Honest, Open, Proud toolkit provides a tool for school professionals to assist students to understand their story from a strength-based perspective and make strategic decisions about sharing highly sensitive and personal information like their mental health or suicide history.

In-School Suicide Attempts

In the case of an in-school suicide attempt, the physical and mental health and safety of the student are paramount. In these situations:

- First aid will be rendered until professional medical services and/or transportation can be received, following district emergency medical procedures.
- School staff will supervise the student to ensure their safety.
- Staff will move all other students out of the immediate area as soon as possible.

- Staff will request a mental health assessment for the youth as soon as feasibly possible. Since self-harm behaviors are on a continuum of level and urgency, and not all instances of suicidal ideation or behavior warrant hospitalization, a mental health assessment including suicide risk assessment can help determine the best treatment plan and disposition.
- The school employed mental health professional or principal will contact the student's parent or guardian, as described in the Working with Parents section.
- Staff will immediately notify the principal or school suicide prevention coordinator regarding the incident of in-school suicide attempt.
- The school will engage their building/district level crisis team, as necessary, to assess whether additional steps should be taken to ensure the safety and well-being of the student, and those who may have had emotional or physical proximity to the victim.

Out-of-School Suicide Attempts

If a staff member becomes aware of a suicide attempt by a student that is in progress in an out-of-school location, the staff member will:

- Call 911 (police and/or emergency medical services).
- Inform the student's parent or guardian.
- Inform supervisor and principal.
- Maintain contact with student if possible until help arrives.

**Adapted from Model School District Policy on Suicide Prevention, AFSP, 2014*

TIER III CONSIDERATIONS

- Student support services staff have professional development in suicide intervention, including the use of brief evidence-based assessment tools and safety planning.
- Procedure for helping a student at risk for suicide is standardized, and includes screening risk assessment, parental notification, brief intervention including safety planning, means restriction, and crisis contacts, referrals, and documentation.
<https://www.ncbi.nlm.nih.gov/pubmed/20634440>
- Student is supervised at all times throughout the assessment and intervention process.
- Principal or designee is notified of intervention as soon as logistically possible.
- Procedure emphasizes collaborative decision-making and support.
- Standardized evidence-based assessment tool is used, and health and mental health staff receive training in its use.
- Assessment of and outreach to community mental health providers to create a network and discuss care collaboration.
- Documentation of suicide risk is standardized and kept secure, and the information is available only to those who need to know it, including the student and their family.
- A policy exists on when and how to notify parents if suicide risk is a concern, and what to do if parents cannot be contacted.
- Parents are notified if a suicide risk assessment is done or if a student is deemed to be at risk for suicide.

Postvention

The death of a student or staff member present challenges to the school community. When the death is by suicide there can be even greater challenges. The complex nature of suicide grief, the sudden, unexpected, and often violent aspects of this kind of loss, and the difficulty we have understanding and talking about suicide create the circumstances that can leave school leaders in shock and struggling to meet the needs of staff and students.


Written protocols, policy, preplanning, well-defined roles, and professional development specific to suicide loss are essential supports that enable school leaders and crisis team members to respond in a coordinated and effective manner. **After a Suicide: A Toolkit for Schools**, published in 2018 by the American Foundation for Suicide Prevention (AFSP) and the Suicide Prevention Resource Center (SPRC), can be used in the immediate aftermath of a suicide death.

Chapter 3 in **Preventing Suicide: A Toolkit for High Schools** has information and templates that can be helpful for policy development and preplanning. Additionally, AFSP, The Trevor Project, and National Association of School Psychologists (NASP) document **Model School District Policy for Suicide Prevention** has a section devoted to suicide postvention guidance.

The Suicide Prevention Center of New York has resource staff who are connected to local resources across NYS that can support schools with short-term and long-term recovery needs through direct consultation and telephone support, or professional development workshops. They support crisis team development through consultation and training based on the **Lifelines Postvention: Responding to Suicide and Other Traumatic Death** model (Underwood, 2017). The goals of postvention should be to resolve the crisis and restore the learning environment, provide appropriate information about the crisis, identify students and staff who may be affected and assure grief and trauma supports are in place, and identify those who may be at risk. A word about **suicide contagion**. Research has demonstrated that exposure to suicide within one's family, in one's peer group, school community, or even through medial sources, can result in increased suicidal behavior and death in others. Direct or indirect exposure to suicidal behavior or suicide death has shown to increase suicide risk in others, especially adolescents and young adults. This risk can be minimized by efforts to provide factual and concise information, offering access to grief and crisis supports, and by checking in with students and staff members who may be vulnerable. (HHS, 2019)

POSTVENTION CONSIDERATIONS

- Develop written procedures and protocols and conduct professional development for crisis team in advance of suicide death in the district.
- Preplanning, well-defined roles, channels of communication, and team collaboration are essential to enable school leadership and crisis team members to respond in a coordinated and effective manner.
- Procedure should be in place for identifying and following up with those affected by a death by suicide. Follow up may include brief group psychoeducational session, grief support, mental health or trauma referrals, or suicide risk screening.
- Care should be taken in communicating information about the death. Verify the facts, avoid speculation, offer helpful resources. Use care not to sensationalize the event, to reduce the risk of contagion.
- Certain memorial activities have been linked to suicide contagion. School leaders should ensure that they are informed about the risks and consider rules that might steer students and staff toward more helpful memorial activities.



Postvention: Immediate and long-term response after suicide loss in MTSS Framework

Tier III – Individualized Supports

Efforts made to quickly identify students who may be directly or indirectly affected. And identify some that may require continued monitoring, care, and referral for grief and/or trauma supports.

Tier II – Targeted Group Supports

Provide small psychoeducational support groups/spaces. Groups may be self-selected or selected by response team. For example, as sports team or close friends of the deceased.

Tier I – Universal Supports

All students receive general announcement about the death, have an opportunity to ask questions, dispel rumors, and learn about funeral arrangements. Often done in small or classroom groups and may include parents.

**Adapted from Erlacher, Singer & Poland 2014.*

Resources

For School Leaders

- **School Readiness Workbook:** This workbook offers a framework, checklists, and resources that school leaders and/or school implementation teams, can use to:
 1. Assess current suicide prevention, intervention, and postvention readiness
 2. Obtain professional development and training for staff
 3. Develop suicide intervention and postvention procedures
 4. Produces a plan to implement suicide prevention, intervention, and postvention initiatives
 5. Find high quality resources
 6. Track effectiveness of efforts
 - ▶ <https://www.preventsuicideny.org/wp-content/uploads/2019/06/School-Readiness-Workbook.pdf>
- **Model School District Policy for Suicide Prevention:** This model policy was created through a collaboration between the Trevor Project, the American Foundation for Suicide Prevention, the American School Counselors Association, and the National Association of School Psychologists.
 - ▶ <https://www.thetrevorproject.org/education/model-school-policy/>
- **Preventing Suicide: A Toolkit for High Schools:** Published by SAMHSA, this toolkit provides best practice guidance and many templates and worksheets. It can be downloaded or hard copies can be ordered at no cost.
 - ▶ <https://store.samhsa.gov/system/files/sma12-4669.pdf>
- **After a Suicide: A Toolkit for Schools** can be printed and offers a grab and go guide for responding.
 - ▶ <http://www.sprc.org/sites/default/files/resource-program/AfteraSuicideToolkitforSchools.pdf>

- **Mental health Education Literacy in Schools: Linking to a Continuum of Well-Being.**
A comprehensive guide for New York Schools published by NYSED.
 - ▶ <http://www.nysed.gov/common/nysed/files/programs/curriculum-instruction/continuumofwellbeingguide.pdf>
- **School Mental Health Resource and Training Center** at MHANYS was established to support schools to implement Chapter 390 of the Law of 2016, requiring mental health instruction as part of the K-12 health curricula.
 - ▶ <https://www.mentalhealthdnys.org/>
- **School and Mental Health Partnerships: Improving School and Community Outcomes for Children and Adolescents with Emotional and Behavioral Challenges.**
 - ▶ <https://www.omh.ny.gov/omhweb/childservice/docs/school-based-mhservices.pdf>
- **Promoting Parent Engagement: Improving Student Health and Academic Achievement**
 - ▶ https://www.cdc.gov/healthyouth/protective/pdf/parentengagement_administrators.pdf

Online Resources For Parents

- **Society for the Prevention of Teen Suicide Parent Pages**
 - ▶ <https://www.sptsusa.org/parents/>
- **Mayo Clinic parent awareness video, brief, warning signs**
 - ▶ <https://www.youtube.com/watch?v=3BBYqa7bhto>
- **The American Academy of Child and Adolescent Psychiatry Suicide Resource Center**
 - ▶ https://www.aacap.org/AACAP/Families_and_Youth/Resource_Centers/Suicide_Resource_Center/Home.aspx

Printable Resources For Parents

- **SPC-NY Parent brochure**
 - ▶ <https://www.preventsuicideny.org/wp-content/uploads/2019/02/ML6-What-every-parent-should-know.pdf>
- **CDC Adolescent and School Health Division** offers School Connectedness for Parents and Families, a printable guide.
 - ▶ https://www.cdc.gov/healthyyouth/protective/pdf/connectedness_parents.pdf
- **Ways to Engage in Your Child's School to Support Student Health and Learning**
 - ▶ https://www.cdc.gov/healthyyouth/protective/pdf/parentengagement_parents.pdf

Online Resources For Educators

- **Act on FACTS: Making Educators Partners in Teen Suicide Prevention** – two-hour online training for educators.
 - ▶ <http://www.sptsusa.org/educators/meptrailer/>
- Two brief videos provide common warning signs of suicide and how to make a warm handoff to school mental health resources.
 - ▶ <https://vimeo.com/272856590>
 - ▶ <https://vimeo.com/272856564>
- Self-care and compassion resilience are two important concepts in protecting the mental health of educators.
 - ▶ <https://rogersbh.org/Insight/teaching-compassion-resilience>

Printable Resources For Educators

- **SPC-NY Teachers brochure** with common warning signs.
 - ▶ https://docs.wixstatic.com/ugd/f5da1_176748adb283415989c01ff3d26d9f0.pdf
- **The role of high school teachers in preventing suicide**
 - ▶ http://www.sprc.org/sites/default/files/resource-program/Role%20of%20High%20School%20Teachers%20Revised%20FINAL%20v2_6-14-19.pdf
- **Wise Wisconsin** offers 7 promises of a trusted adult and a Safe Persons Decal for those who challenge themselves to uphold the 7 promises.
 - ▶ <https://wisewisconsin.org/blog/the-safe-person-decal/>

Online Resources For Working with Students

- **27 Resiliency Activities and Worksheets for Students and Adults**
 - ▶ <https://positivepsychologyprogram.com/resilience-activities-worksheets/#schools>
- **Sources of Strength:** This website not only offers information about how schools can enroll in the Sources of Strength program, it also offers plenty of resources and tips about how students can run positive, strength-based messaging campaign.
 - ▶ <https://sourcesofstrength.org/peer-leaders/>
- **Intervention Central** offers information on how to implement the Good Behavior Game, an elementary classroom intervention and evidence-based practice for improving pro-social behavior.
 - ▶ <http://www.interventioncentral.org/behavioral-interventions/schoolwide-classroommgmt/good-behavior-game>
- **DBT STEPS A:** a universal intervention teaching the skills to manage emotional distress and targeting emotional dysregulation which has been implicated in suicidal ideation and self-injury.
 - ▶ <http://www.dbtinschools.com>
- **Strategic Disclosure workbook and facilitators guide** helps students with mental health challenges gain a sense of personal empowerment and confidence in achieving goals through reducing self-stigma.
 - ▶ <https://wisewisconsin.org/honest-open-proud/>
- This program offers 10 free lesson plans for use in classrooms to teach the skills to find and maintain hope.
 - ▶ <https://www.teachthought.com/life/preventing-youth-suicide-through-lessons-of-hope/>
- **National Center for Safe and Supportive Learning Environments** offers free School Climate Surveys.
 - ▶ **School Climate survey and resources**
- **Social Emotional Learning: A guide to Systemic Whole School Implementation.**
 - ▶ <http://www.p12.nysed.gov/sss/documents/SystemicWholeSchoolImplementationFINAL.pdf>

- ▶ **SPTSUSA.org/Teens – Society for the Prevention of Teen Suicide**
- **The Trevor Project** offers telephone, text and chat support and crisis services for LGBTQ+ youth. They also offer ally training to adults who support youth.
 - ▶ <https://www.thetrevorproject.org/get-help-now/>
- This video created by the **Mayo Clinic** encourages teens to reach out to trusted adults for help.
 - ▶ <https://www.youtube.com/watch?v=TleBxeOxFyE>

Printable Resources For Working with Students

- **SPC-NY Awareness brochure**
 - ▶ <https://www.preventsuicideny.org/wp-content/uploads/2019/02/ML5-Be-a-good-Friend.pdf>
- **CDC Adolescent and School Health Division - Fostering School Connectedness: Information for Teachers and Other School Staff**
 - ▶ https://www.cdc.gov/healthyouth/protective/pdf/connectedness_teachers.pdf
- **SPC-NY Talking Points for classroom discussions on suicide.**
 - ▶ https://docs.wixstatic.com/ugd/da1_2a35f0f6e25148bc97b3a1596321abc5.pdf

Online Resources For School-based Health and Mental Health Professionals

- **The Columbia Lighthouse Project** offers training and free access to protocols and suicide assessment and screening tools, customized for a variety of settings including schools.
▶ <http://cssrs.columbia.edu/>
- **Center for Practice Innovations at New York State Psychiatric Institute** offers free online training in the AIM-SP model for Suicide Safer Care for Children and Adolescents.
▶ <https://practiceinnovations.org/i-want-to-learn-about/Suicide-Prevention/Trainings/Full-list-of-SP-TIE-trainings#top>
- **Safety Planning Intervention template with telephone app.**
▶ <https://my3app.org/>
- **Safety Plan Template for use on electronic devices.** Safety planning is an evidence-based intervention for suicidal individuals.
▶ <https://itunes.apple.com/us/app/suicide-safety-plan/id1003891579?mt=8>
- **NYS Office of Mental Health School and Mental Health Partnerships:** What School Leaders Should Know When Creating School and Mental Health Partnerships.
▶ <https://www.omh.ny.gov/omhweb/childservice/docs/mh-101-for-educators.pdf>
- **A Practitioner's Resource Guide: Helping Families Support Their LGBT Children.**
▶ <https://familyproject.sfsu.edu/sites/default/files/FamilySupportForLGBTChildrenGuidance.pdf>
- **The Role of High School Mental Health Providers in Suicide Prevention**
▶ http://www.sprc.org/sites/default/files/resource-program/Role%20of%20HS%20MH%20Providers%20Revised%20FINAL%20v3_6-18-19.pdf

Printable Resources For School-based Health and Mental Health Professionals

- **SPC-NY Awareness brochure**
▶ <https://www.preventsuicideny.org/wp-content/uploads/2019/02/ML5-Be-a-good-Friend.pdf>
- **CDC Adolescent and School Health Division - Fostering School Connectedness: Information for Teachers and Other School Staff**
▶ https://www.cdc.gov/healthyyouth/protective/pdf/connectedness_teachers.pdf
- **SPC-NY Talking Points for classroom discussions on suicide.**
▶ https://docs.wixstatic.com/ugd/da1_2a35f0f6e25148bc97b3a1596321abc5.pdf

References


- American Foundation for Suicide Prevention, The Trevor Project, National Association of School Psychologists, American School Counselor Association. (2014). Model School District Policy on Suicide Prevention: Model Language, Commentary, and Resources. Retrieved from https://afsp.org/wp-content/uploads/2016/01/Model-Policy_FINAL.pdf
- Barnes, A. J., Gilbertson, J., & Chatterjee, D. (2018). Emotional Health Among Youth Experiencing Family Homelessness. *Pediatrics*,141(4). doi:10.1542/peds.2017-1767
- Becerra-Culqui, T. A., Liu, Y., Nash, R., Cromwell, L., Flanders, W. D., Getahun, Giammattei, S. V., Hunkeler, E. M., Lash, T. L., Millman, A., Quinn, V. P. Robinson, B., Roblin, D., Sandberg, D. E., Silverberg, M. J., Tangpricha, V., Goodman, M. (2018). Mental Health of Transgender and Gender Nonconforming Youth Compared with Their Peers. *Pediatrics*, Volume 141, Issue 5. Retrieved from <https://pediatrics.aappublications.org/content/141/5/e20173845>
- Breux, P., Boccio, D., Brodsky, B. (2017). Creating Suicide Safety in Schools: A public health suicide prevention program in New York State. *Suicidologi*, Vol 22, Nr 2. Pg 14 – 26.
- Center for Disease Control and Prevention. (2018). Youth Risk Behavior Survey Data Summary and Trends Report 2007 – 2017. <https://www.cdc.gov/healthyyouth/data/yrbs/pdf/trendsreport.pdf>
- Center for Disease Control and Prevention. (2014). The Relationship Between Bullying and Suicide: What We Know and What it Means for Schools. Retrieved from <https://www.cdc.gov/violenceprevention/pdf/bullying-suicide-translation-final-a.pdf>
- Center for Disease Control and Prevention. (2019). Fatal Injury Reports, National, Regional and State, 1981-2017. Retrieved from <https://webappa.cdc.gov/sasweb/ncipc/mortrate.html>
- Center for Disease Control and Prevention. (2018). Data and Statistics (WISQARS). 10 Leading Causes of Death by Age Group, United States - 2017. Retrieved from https://www.cdc.gov/injury/images/lc-charts/leading_causes_of_death_by_age_group_2017_1100w850h.jpg
- Center for Disease Control and Prevention. (2019). Violence Prevention. The Social-Ecological Model: A Framework for Prevention. Retrieved from <https://www.cdc.gov/violenceprevention/publichealthissue/social-ecologicalmodel.html>
- Center for Disease Control and Prevention. (2014). The Relationship Between Bullying and Suicide: What We Know and What It Means for Schools.
- Center for Disease Control and Prevention. (2018). High School Youth Risk Behavior Survey New York Results 2017. Retrieved from <https://nccd.cdc.gov/youthonline/App/Results.aspx?LID=NY>
- Erbacher, T., Singer, J., Poland, S. (2015). *Suicide in Schools: A Practitioner's Guide to Multi-level Prevention, Assessment, Intervention, and Postvention*. Rutledge, New York.
- Guldin, M., Li, J., Pedersen, H. S. (2015). Incidence of Suicide Among Persons Who Had a Parent Who Died During Their Childhood: A Population-Based Cohort Study. *JAMA Psychiatry*. 2015;72(12);1227-1234. DOI:10.1001/jamapsychiatry.2015.2094
- Hatzenbuehler, M. I. (2011). The Social Environment and Suicide Attempts in Lesbian, Gay, and Bisexual Youth. *Pediatrics* 2011 May; 127 (5): 896-903. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3081186/>
- Humenski, J., Coronel, B., Mazzula, S., Lewis-Fernandez, R. (2016). Life is Precious: A Community-Based Program to Reduce Suicidal Behavior in Latina Adolescents. *Archives of Suicide Research*. DOI: 10.1080/13811118.2016.1242442

- Jones, S.E., Lollar, D. J. (2008). Relationship Between Physical Disabilities or Long-Term Health Problems and Health Risk Behaviors or Conditions Among US High School Students. *Journal of School Health* Volume 78, Issue 5 May, 2008. Pages 252-257. Retrieved from <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1746-1561.2008.00297.x>
- King, C., Foster, C., Rogalski, K. (2013). *Teen Suicide Risk*. Guilford, London. Pg. 14-19, 33,34.
- Klonsky, D. E., Victor, S. E. Saffer, B. Y. (2014). Nonsuicidal Self-Injury: What We Know, And What We Need to Know. *Canadian Journal of Psychiatry*, 2014 Nov; 59(11): 565-568 Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4244874/>
- Kutcher, S., Wei, Y., Coniglio, C. (2016) Mental Health Literacy Past, Present, and Future. *Canadian Journal of Psychiatry*. 2016 Mar; 61 (3): 154-158. Doi: 10.1177/0706743715616609. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4813415/>
- Ludi, E., Ballard, E.D., Greenbau, R., Pao, M., Bridge, J., Reynolds, W., Horowitz, L. (2013). Suicide Risk in Youth with Intellectual Disability: the Challenges of Screening. *Journal of Developmental and Behavioral Pediatrics*. 2012 June; 33(5): 431-440. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3464013/>
- Mental Health Technology Transfer Center Network. (2019). *After a School Tragedy... Readiness, Response, Recovery, & Resources*. Retrieved from <https://mhttcnetwork.org/centers/mhrtc-network-coordinating-office/after-school-tragedyreadiness-response-recovery-resources>
- New York State Department of Health. (2018). *Toward the Healthiest State Progress Report 2018. Contributing Causes of Health Challenges*. https://health.ny.gov/prevention/prevention_agenda/2019-2024/docs/sha/contributing_causes_of_health_challenges.pdf
- New York State Education Department. (2015). *Guidance to School Districts for Creating a Safe and Supportive School Environment for Transgender and Gender Nonconforming Students*. Retrieved from http://www.p12.nysed.gov/dignityact/documents/Transg_GNCGuidanceFINAL.pdf
- New York State Office of Mental Health. (2016). *1,700 Too Many: New York State's Suicide Prevention Plan 2016-2017*.
- Nickerson, A., Singleton, D., Schnurr, B., Collen, M.H. (2014). Perceptions of School Climate as a function of Bullying Involvement. *Journal of Applied School Psychology*. Vol. 30, 2014 – Issue 2. P 157 – 181.
- Okpych, N. J., & Courtney, M. E. (2018). Characteristics of Foster Care History as Risk Factors for Psychiatric Disorders Among Youth in Care. *American Journal of Orthopsychiatry*, 88(3), 269–281. doi:10.1037/ort0000259
- Parra-Uribe, I., Blasco-Fontecilla, H., Garcia-Parés, G., Martínez-Naval, L., Valero-Coppin, O., Cebrià-Meca, A., ... Palao-Vidal, D. (2017). Risk of re-attempts and suicide death after a suicide attempt: A survival analysis. *BMC Psychiatry*, 17(1), 163. doi:10.1186/s12888-017-1317-z
- Scott, M., Underwood, M., & Lamis, D. (2015). Suicide and Related-Behavior Among Youth Involved in the Juvenile Justice System. *Child & Adolescent Social Work Journal*, 32(6), 517–527. doi: 10.1007/s10560-015-0390-8
- Robinson, J.P., Espelage, D. L. (2012). Bullying Explains Only Part of LGBTQ-Heterosexual Risk Disparities: Implications for Policy and Practice. *American Educational Research Association*. Volume: 41 Issue: 8. Pages: 309-319. Retrieved from <https://journals.sagepub.com/doi/10.3102/0013189X12457023>
- Sandler, I., Tein, J.Y., Wolchik, S. A., Ayers, T. (2016). The effects of the Family Bereavement Program to reduce suicide ideation and/or attempts of parentally bereaved children six to fifteen years later. *Suicide and Life Threatening Behavior*, 46, S32-S38. DOI: 10.1111/sltb. 12256.


- Schaffer, G. (2017). Nuts & Bolts: Multi-Tiered Systems of Support: A Basic Guide to Implementing Preventative Practices in Our Schools. New York Association of School Psychologists. Albany.
- Stanley, B., Brown, G. K. (2012). Safety Planning Intervention: A Brief Intervention to Mitigate Suicide Risk. Cognitive and Behavioral Practice. 19 (2012) 256-264. Retrieved from http://www.suicidesafetyplan.com/uploads/Safety_Planning_-_Cog__Beh_Practice.pdf
- Substance Abuse and Mental Health Services Administration. (2012) Preventing Suicide: A Toolkit for High Schools. HHS Publication No. SMA-12-4669. Rockville, MD: Center for Mental Health Services.
- Substance Abuse and Mental Health Services Administration. (2016). Substance Use and Suicide: A Nexus Requiring a Public Health Approach. In Brief. Retrieved from <https://store.samhsa.gov/system/files/sma16-4935.pdf>
- Suicide Prevention Center of New York. (2017). Suicide Safety for Teachers and School Staff Trainer Manual.
- U. S. Department of Health & Human Services. (2019). What does "suicide contagion" mean, and what can be done to prevent it? Retrieved from <https://www.hhs.gov/answers/mental-health-and-substance-abuse/what-does-suicide-contagion-mean/index.html>
- Underwood, M. (2010). Creating Competent Communities for Youth Suicide Prevention. Powerpoint presentation. Society for the Prevention of Teen Suicide. <https://1.cdn.edl.io/P12Qlrci9ugxvZ7sBkVvOELwfhgdZxTlsgkQLxS.pdf> • <https://www.cdc.gov/healthyschools/wsc/index.htm>
- Underwood, M. (2018). Suicide Prevention in Schools, Issue Brief. Substance Abuse and Mental Health Services Administration. Project Aware. Retrieved from <http://files.constantcontact.com/bde05f96001/5641d914-e6d6-439b-b7c6-e3f4ca09d61f.pdf>
- Underwood, M., Fell, F., Spinazzola, N. (2018). Lifelines Postvention: Responding to Suicide and Other Traumatic Death. Hazelden, Center City.
- World Health Organization. (2014). Preventing Suicide A Global Imperative. Retrieved from https://apps.who.int/iris/bitstream/handle/10665/131056/9789241564779_eng_FFE81908E9580A06CA24384B?sequence=1
- World Health Organization. (2002). Violence-a global public health problem. In: Krug, E., Dahlberg, L.L., Mercy, J.A., Zwi, A.B., Lozano, R., eds. World Report on Violence and Health. Geneva, Switzerland. Retrieved from https://www.who.int/violence_injury_prevention/violence/world_report/en/chap1.pdf
- Wyman, P.A., Brown, C.H., LoMurray, M., Schmeelk-Cone, K., Petrova, M., Yu, Q., Walsh, E., Tu, X., Wang, W. (2010). An outcome evaluation of the Sources of Strength suicide prevention program delivered by adolescent peer leaders in high schools. American Journal of Public Health. 2010 September 100(9); 1653-61. Doi: 10.2105/AJPH.2009.190025. Epub 2010 Jul 15. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/20634440>
- Youth Risk Behavior Survey Data Summary & Trends Report 2007 to 2017. (2018) National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, Division of Adolescent and School Health. Retrieved from <https://www.cdc.gov/healthyyouth/data/yrbs/pdf/trendsreport.pdf>
- Zayas, L. H., Pilat, A. M. (2009). Suicidal Behavior in Latinas: Explanatory Cultural Factors and Implications for Intervention. Suicide and Life-Threatening Behavior. 38(3) 334-342. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2662359/#R8>
- Zimmerman, M.A. (2013). Resiliency Theory: A Strengths-Based Approach to Research and Practice for Adolescent Health. Health Education & Behavior. <https://doi.org/10.1177%2F1090198113493782>



American Foundation for Suicide Prevention



jedfoundation.org



YOUTH POWER!

IF YOU NOTICE ANY OF THESE WARNING SIGNS, TAKE ACTION!

Signs that a student may be at risk include the following F-A-C-T-S:

- **FEELINGS** like expressing hopelessness about the future, seeming sad and unhappy, being anxious and worried, or getting angry and aggressive.
- **ACTIONS** like withdrawing from activities or friendships, doing risky, dangerous things like drinking & driving, or researching ways to die online.
- **CHANGES** in the normal mood and behavior of your student. In some ways, this may be what is easiest for you to notice. If you observe changes that concern you, reach out to others in the student's life (i.e., parents, teachers, friends, religious leaders, etc.) to see if they've also noticed changes.
- **THREATS** are sometimes direct like "I'd rather be dead". They can also be vague like "I just don't care about anything anymore."
- **SITUATIONS** are events that can serve as triggers for the suicidal behavior. These can include things like getting into trouble at home or school or with the law, experiencing some type of loss or facing a life change that may be too overwhelming for the student to deal with on their own.

SUICIDE IS PREVENTABLE.

By taking time to notice and reach out to someone you feel is at risk, **you** can be the beginning of a positive solution.



Office of
Mental Health


WHAT EVERY TEACHER NEEDS TO KNOW:

Recognizing Suicide Risk in Students



CRISIS TEXT LINE |

Text "START" to 741-741



www.PreventSuicideNY.org

**Youth Suicide Prevention
is Everyone's Business!**

YOUR ROLE AS A TEACHER IS CRITICAL

Does teaching seem to get harder every year?

Are there more requirements, more testing, and less time for you to think— let alone plan?

Are you expected to take more responsibilities for your students, even when they are more challenging and when some of them may be at-risk for suicide?

Did you know that according to national data:

Almost 30% of 9th through 12th grade students have felt so sad or helpless during the course of an academic year that they couldn't do the things they normally do?

Or that there has been a dramatic rise in the suicide attempt rate for 10 to 14 year olds?

Or that suicide is the 2nd leading cause of death for youth in New York State?



Who are these kids?

They're sitting in your classrooms every day.

Although your job is to teach them, not diagnose them, there are ways that can help you better identify these struggling students and get them to someone who is trained to make a more complete assessment of their needs. Students who are thinking about suicide are not concentrating on school work; they are often preoccupied with problems that seem overwhelming and unsolvable.

Your role in this process is critical but very limited and is often the first step in getting students the help they need.

So how do you accomplish this?

By doing what you do best- simply paying attention to your students and knowing where to send them in your school if you notice anything that concerns you.

The majority of those students who are thinking about suicide show direct or indirect warning signs. These are things that reflect a change in the student's behavior, attitude or feelings from as little as two weeks ago.

Some common warning signs are listed on the back panel of this brochure. If you see any of these, your responsibility is to get that student to the appropriate resources in your building.

Remember, your job isn't to figure out what the problem is— it's simply to get this student help.

Be sure to follow up with that resource person to ensure action is being taken and check in with the student to see how things are going. If you continue to be concerned, let that resource person know.

Suicide risk doesn't immediately disappear once an intervention is made, so keep your eyes open!

Noticing and referring potentially at-risk students are only the beginning of the suicide prevention equation. Equally important is your role in encouraging students to seek help if they have a problem and to turn to a trusted adult for support.

Help-seeking is called a protective factor, the kind of thing that can buffer us from life stressors.

The single most important protective factor for youth is a relationship with one trusted adult. As you know too well, many of your students may not have very supportive situations outside of school, so their trusted adult is often someone in their school community.

What does it take to be a trusted adult to a student?

Here's how students describe it:

- Making time to talk, even if your schedule is tight
- Taking my concerns seriously, no matter how trivial they seem
- Not telling me “ it will be better tomorrow”
- LISTENING! Recognizing you probably can't fix what I'm worried about but just listening to me talk about it can help
- Being honest if you think you have to tell someone else about my problem
- Taking action when it's necessary
- Remembering what we talked about and asking me about it later

When you review this list, you'll probably find that these are the same things you look for in someone to whom you turn for help— it's no different! While simply listening to a student talk about suicide can be very difficult, remember, it's the first step in the process.

That critical next step is getting that student to the resources in your school that can offer more help!